



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 610 511 A1**

(12)

EUROPEAN PATENT APPLICATION
published in accordance with Art.
158(3) EPC

(21) Application number: **93914996.9**

(22) Date of filing: **13.07.93**

(86) International application number:
PCT/JP93/00968

(87) International publication number:
WO 94/01083 (20.01.94 94/03)

(51) Int. Cl.⁵: **A61K 7/48, A61K 7/00,
A61K 31/57, A61K 31/19**

(30) Priority: **13.07.92 JP 227723/92**
13.07.92 JP 227724/92
13.07.92 JP 227726/92

(43) Date of publication of application:
17.08.94 Bulletin 94/33

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC
NL PT SE**

(71) Applicant: **SHISEIDO COMPANY LIMITED**
7-5-5 Ginza
Chuo-ku Tokyo 104 (JP)

(72) Inventor: **YANAGIDA, Takeshi The Shiseido**
Research Center
1050, Nippa-cho
Kohoku-ku
Yokohama-shi Kanagawa 223 (JP)
Inventor: **SAKAMOTO, Okihiko The Shiseido**
Research Center
1050, Nippa-cho
Kohoku-ku
Yokohama-shi Kanagawa 223 (JP)

(74) Representative: **Rinuy, Santarelli**
14, avenue de la Grande Armée
F-75017 Paris (FR)

(54) **COMPOSITION FOR DERMATOLOGIC PREPARATION.**

(57) A composition for dermatologic preparation comprising vitamin A and at least one chap ameliorant aid selected from the group consisting of polyoxyalkylene-modified organopolysiloxane, sugars and anti-inflammatory.

EP 0 610 511 A1

TECHNICAL FIELD

The present invention relates to an external skin treatment composition and, more specifically, it relates to an external skin treatment composition having a synergistically improved skin roughening improvement effect by incorporating therein vitamin A and a polyoxyalkylene modified organopolysiloxane, a sugar or an anti-inflammatory agent, with taking safety into consideration.

BACKGROUND ART

Various pharmaceutically effective components are formulated into external skin treatment compositions. Among these pharmaceutical effects, an effect, by which the changes in the skin due to aged skin or sunlight exposure etc. are prevented or improved, is one of such effects, and therefore, an external skin treatment compositions such as cosmetic compositions having such purposes have been desired.

Under such circumstances, various raw materials extracted from natural products, such as proteins, polysaccharides, extracted extracts, natural polymers etc. have been heretofore formulated in external skin treatment compositions due to their characteristics application effects.

Recently, Japanese Unexamined Patent Publication (Kokai) No. 64-500355 discloses a method for preventing or improving the changes or disabilities caused due to aged skins or sunlight exposure by formulating therein at least one component selected from vitamin A and the derivatives thereof.

However, the effects thereof are not sufficient and it has been strongly desired to develop a pharmaceutically effective agent having much more excellent effects.

DISCLOSURE OF THE INVENTION

Accordingly, the objects of the present invention are to obviate the above-mentioned problems in the prior art and to provide an external skin treatment composition having sufficient skin roughening improvement effects, i.e., prevention effects, improvement effects, etc. against the changes or disabilities due to aged skin or sunlight exposure.

In accordance with the present invention, there is provided an external skin treatment composition comprising (i) vitamin A and (ii) at least one skin roughening improvement aid selected from the group consisting of (a) polyoxyalkylene modified organopolysiloxanes, (b) sugars, and (c) anti-inflammatory agents.

Best Mode for Carrying out the Invention


In order to achieve the above-mentioned objects, the present inventors have been extensively studied to obtain a substance or substances capable of effecting sufficient skin roughening improvement effects, especially, among substances having excellent safety and, as a result, found that the above-mentioned problems can be solved by formulating, together with vitamin A, polyoxyalkylene modified organopolysiloxanes, sugars or anti-inflammatory agents.

The constitution of the present invention will now be explained in detail.

Vitamin A used in the present invention is also called retinol and is usually used in the treatment of infant or childhood diseases or nyctalopia (i.e., night blindness) or in the recovery agent after pregnancy in the pharmaceutical fields etc. Among these, all-trans products or 13-cis products can be preferably used, but the mixture thereof can also be used.

There are no limitations to the amounts of vitamin A formulated into the external skin treatment agent according to the present invention, but the preferable amount is 0.00001 to 5% by weight, more preferably 0.0001 to 0.5% by weight, in view of the effect of vitamin A to the skin.

The polyoxyalkylene modified organopolysiloxanes usable as a skin roughening improvement aid, in the present invention are the following compounds (A), (B), (C) and (D).



wherein R represents an alkyl group having 1 to 3 carbon atoms, or a phenyl group, R' represents hydrogen or an alkyl group having 1 to 12 carbon atoms, p is an integer of 1 to 5, m is an integer of 5 to 100, n and x are an integer of 1 to 50 and t and y are an integer of 0 to 50.

There are no specific limitations to the average molecular weight of the polyoxyalkylene modified organopolysiloxane usable in the present invention, but the preferable molecular weight is 3,000 or more,

further preferably 5,000 to 10,000. Furthermore, the preferable polyoxyalkylene modified organopolysiloxanes are those having 2 - 80% by weight, more preferable, 11 - 50% by weight in view of the generation of the effects, of polyoxyalkylene group in the molecule thereof.

The amount of the polyoxyalkylene modified organopolysiloxane formulated is preferably 0.1 to 20% by weight, more preferably 0.2 to 10% by weight, in the total amount of the external skin treatment composition. When the amount is less than 0.1% by weight, there are fears that the skin irritation is not sufficiently lowered. Contrary to this, when the amount is more than 20% by weight, there are fears that the qualities necessary as the skin treatment composition cannot be held.

In the present invention, as the sugars usable as a skin roughening improvement aid, mention may be made of monosaccharides, oligosaccharides, sugar alcohols, etc.

As the monosaccharides, mention may be made of trioses such as D-glyceraldehyde, dihydroxyacetone, etc., tetroses such as D-erythrose, D-erythritolose, D-threose, etc., pentoses such as L-arabinose, D-xylose, L-lyxose, D-arabinose, D-ribose, D-ribulose, D-xylulose, L-xylulose, etc., hexoses such as D-glucose, D-talose, D-psicose, D-galactose, D-fructose, L-galactose, L-sorbose, D-mannose, D-tagalose etc., heptoses such as aldoheptose, heptalose etc., octoses such as octose etc., deoxy sugars such as 2-deoxy-D-ribose, 6-deoxy-L-galactose, 6-deoxy-L-mannose, etc., amino sugars such as D-glucosamine, D-galactosamine, sialic acid, aminouronic acid, muramic acid, etc., uronic acids such as D-glucuronic acid, D-mannuronic acid, L-gluconic acid, D-galacturonic acid, L-iduronic acid etc.

As the oligo sugars, mention may be made of sucrose, gentianose, umbelliferose, lactose, planteose, α,α -trehalose, raffinose, umbilicin, stachyose, verbascoses.

Furthermore, as the sugar alcohols, mention may be made of sorbitol, maltiol, maltriose, mannitol, starch decomposed sugar, erythritol, xylitose, starch decomposing sugar reduced alcohols. Of these sugar alcohols, mannitol, erythritol and sugar alcohols of disaccharides or more.

There are no specific limitations to the amount of the sugars formulated, the preferable amount is at least 0.1% by weight, more preferably 0.5 to 50% by weight, based on the total amount of the external skin treatment agent. When the amount is less than 0.1% by weight, it is not preferable because it is difficult to obtain the synergistical skin roughening improvement effects according to the present invention.

Examples of anti-inflammatory agents usable as the skin roughening improvement aid according to the present invention are hydrocortisone, hydrocortisone acetate, prednisolone, methyl prednisolone, prednisolone acetate, prednisolone acetate propionate, dexamethasone, betamethasone, triamcinolone, dexamethasone acetate, betamethasone valerate, triamcinolone acetate, aspirin, salicylic acid, acetaminophen, methyl salicylate, glycol salicylate, mefenamic acid, flufenamic acid, indometacin, diclofenac, ketoprofen, ibuprofen, flurbiprofen, fenbufen, lufexamac, piroxicam, oxyphenbutazone, mepirizole, ibuprofen piconol, clidanac, phenylbutazone, naproxen, glycyrrhetin, glycyrrhizin, glycyrrhetic acid and the salts and esters thereof, glycyrrhizic acid and the salts and esters thereof, azulene, camphor, thymol, allantoin, etc. Among these agents, one or more agents may be freely selected. Although the amount of the anti-inflammatory agents formulated in the present invention is not specifically limited, the preferable amount is 0.0001 to 5.0% by weight, more preferably 0.001 to 2% by weight, based upon the total amount of the external skin treatment composition. When the amount is less than 0.0001% by weight, there are fears that the reduction of the skin irritation, which is the effect of the present external skin treatment composition, cannot be achieved. Contrary to this, even when the anti-inflammatory agent is formulated in an amount of more than 5.0% by weight, further improvement is not expected.

In addition to the above-mentioned essential components, the external skin treatment composition according to the present invention may contain various components conventionally formulated into cosmetics, quasi-drugs, drugs such as aqueous components, humectants, thickeners, UV absorbers, antiseptics, antioxidants, flavours, colorants, medicines, crude drugs, in an amount such that the desired effects of the present invention are not impaired. It should be, of course, noted that these additives are used in such quantitative, qualitative conditions that the objects of the present invention are not impaired.

The external skin treatment composition according to the present invention can be in any form, for example, in the form of a solubilized type such as cosmetic lotions, an emulsified type such as Emulsions, creams, ointments, powder dispersion type, water-oil two layer type, water-oil-powder three layer type, etc.

EXAMPLES

The present invention will now be further illustrated by no means limited to, the following Examples, in which the amounts formulated are "% by weight".

Examples 1-1 to 1-5

Creams having the following compositions were prepared and the skin roughening improvement effects thereof were studied.

5 The polyoxyalkylene modified organopolysiloxanes listed in Table 1-1 were formulated.

		%
10	(1) Cetostearyl alcohol	3.5
	(2) Squalane	30.0
	(3) Beeswax	3.0
	(4) Reduced lanolin	5.0
	(5) Ethylparaben	0.3
	(6) Polyoxyethylene (50 mol) oleyl alcohol ether	2.0
15	(7) Stearic monoglyceride	2.0
	(8) Polyoxyalkylene modified organopolysiloxane (see Table 1-1)	3.0
	(9) Perfume	0.03
	(10) Vitamin A	0.0001
	(11) Glycerol	15.0
20	(12) Purified water	Balance

(Preparation method)

25 (1), (2), (3), (4), (5), (6), (7), (8), (9) and (10) were heated and dissolved. The resultant solution was maintained at 75 °C, followed by adding thereto, under stirring, (11) and (12) heated to 75 °C. The mixture was stirred and emulsified in a homomixer, followed by cooling to obtain a cream.

Comparative Example 1-1

30 The same formulation except that the polyoxyalkylene modified organopolysiloxane was removed from the formulation of Example 1-1.

Comparative Example 1-2

35 The same formulation except that the vitamin A was removed from the formulation of Example 1-1.

Table 1-1

40	No.	Polyoxyalkylene modified organosiloxane
	Example 1-1	General formula A Polyoxyethylene group 20 wt%, M.W. 6,000
45	Example 1-2	General formula A Polyoxyethylene group 40 wt%, M.W. 20,000
	Example 1-3	General formula B Polyoxyethylene group 60 wt%, M.W. 10,000
50	Example 1-4	General formula C Polyoxyethylene group 20 wt%, Polyoxypropylene group 10 wt%, M.W. 4,000
55	Example 1-5	General formula A Polyoxyethylene group 15 wt%, M.W. 2,500

Skin Roughening Improvement Test Method

One hundred subjects having psoriasis-like and skin-roughening-like skin affections were divided into 5 groups as test panels and the creams of Examples 1-1 to 1-5 and Comparative Examples 1-1 to 1-2 were used by one group (20 members) each. That is, Examples 1-1 to 1-5 samples were applied to the left-side faces of the panel twice a day and Comparative Examples 1-1 and 1-2 were applied to the right-side faces of the panel for continuous 3 months. Thereafter, the degree of the overall improvements after the use was visually determined, when compared before the use.

The results are shown in Table 1-2.

10

15

20

25

30

35

40

45

50

55

Table 1-2: Test Results of Practical Use (Degree of Overall Improvement)

Degree of overall improvement	Example					Comparative Example				
	1-1	1-2	1-3	1-4	1-5	1-1	1-1	1-1	1-2	1-2
	Psori- Skin asis rough- ening	Psori- Skin asis rough- ening	Psori- Skin asis rough- ening	Psori- Skin asis rough- ening	Psori- Skin asis rough- ening	Psori- Skin asis rough- ening	Psori- Skin asis rough- ening	Psori- Skin asis rough- ening	Psori- Skin asis rough- ening	Psori- Skin asis rough- ening
Remarkable improvement	7	6	6	6	6	5	5	5	0	0
Some improvement	2	2	1	2	2	3	3	4	4	1
No change	1	2	3	1	2	2	2	19	19	24
Change for the worse	0	0	0	0	0	0	0	2	2	0
Total (No. of Person)	10	10	10	10	10	10	10	25	25	25
Degree of effectiveness %										
Those better than "some improvement"	90	80	90	80	80	80	80	16	16	4

As is clear from the results shown in Table 1-2, the products of Examples 1-1 to 1-5 according to the present invention had synergistically excellent skin roughening improvement effects when compared with those of Comparative Examples 1-1 and 1-2.

Example 1-6: Cosmetic lotion

		%
5	(1) Vitamin A	0.00001
	(2) Polyoxyalkylene modified organopolysiloxane ^{*1}	0.1
	(3) Glycerol	1.0
	(4) D-mannitol	0.5
10	(5) Purified water	Balance
	(6) Ethanol	7.0
	(7) Polyoxyethylene (50 mol) oleyl alcohol ether	1.0
	(8) Methylparaben	0.05
	(9) Oleyl alcohol	1.0
15	(10) Lactic acid	0.01
	(11) Sodium lactate	0.1
	(12) Perfume	0.01

^{*1}: General formula B (Polyoxyethylene group 60 wt%, M.W. 10000)

20

(Preparation method)

In the purified water, (3), (10) and (11) were dissolved. Separately, (1), (2), (7), (8) and (12) were dissolved in ethanol and this solution was added to the above purified water to be dissolved, followed by filtration. Thus, the cosmetic lotion was obtained. The cosmetic lotion of the present invention was excellent in the skin roughening improvement effects.

30

Example 1-7: Pack

35

40

45

		%
	(1) Polyoxyalkylene modified organopolysiloxane ^{*2}	3.0
	(2) Polyvinyl alcohol	10.0
	(3) Propylene glycol	7.0
	(4) Ethanol	10.0
	(5) Vitamin A	0.01
	(6) Methylparaben	0.05
	(7) POE(60 mol) hydrogenated castor oil	0.2
	(8) Perfume	0.05
	(9) Purified water	Balance

^{*2}: General formula A (Polyoxyethylene group 40 wt%, M.W. 8000)

(Preparation method)

(1), (3), (6) and (7) were added to (7) and dissolved under stirring. Then, (2) was added thereto and stirred under heating, followed by adding thereto (4) containing (9) dissolved therein. The mixture was dissolved while stirring to obtain the pack.

The present pack had excellent skin roughening improvement effects.

55

Example 1-8: Compact face powder

		%
5	(1) Vitamin A	0.0005
	(2) Talc	85.4
	(3) Stearic acid	2.5
	(4) Squalane	3.5
10	(5) Sorbitan sesquioleic ester	1.8
	(6) Triethanolamine	1.2
	(7) Polyoxyalkylene modified organopolysiloxane ^{*3}	10.0
	(8) Glycyrrhetic stearyl	0.1
	(9) Pigment	q.s.
15	(10) Perfume	q.s.

^{*3}: General formula B (polyoxyethylene group 60 wt%, M.W. 4000)

20 (Preparation method)

The talc and the pigment were sufficiently mixed by a kneeder (Powder portion). The triethanolamine was added to 50% corresponding amount of the purified water and the mixture was maintained at 70 °C (Aqueous phase). The components of the present invention other than the perfume were mixed and dissolved under heating at 70 °C (Oil phase). The oil phase was added to the aqueous phase, followed by uniformly emulsified by a homomixer and the resultant emulsified mixture was added to the powder portion, followed by kneeding the same by a kneeder, followed by evaporating the water and by treating the same by a grinder. Furthermore, the perfume was uniformly sprayed and the resultant product was compression molded.

30 The resultant compact face powder was excellent in the improvement effects to the skin.

Example 1-9: Lipstick

		%
35	(1) Vitamin A	0.00001
	(2) Microcrystalline wax	3.0
	(3) Beeswax	3.0
40	(4) Ceresin wax	5.0
	(5) Liquid paraffin	19.0
	(6) Squalane	20.0
	(7) Carnauba wax	3.0
	(8) Candelilla wax	3.0
45	(9) Polyoxyalkylene modified organopolysiloxane ^{*4}	1.0
	(10) Mixed colorant	7.0
	(11) Dibutyl hydroxytoluene	0.05
	(12) Perfume	q.s.
50	(13) Lanolin	Balance

^{*4}: General formula A (Polyoxyethylene group 60 wt%, M.W. 25000)

(Preparation method)

55

The lipstick was obtained in a conventional way. The present lipstick exhibited remarkable prevention of the generation of roughening on the lips.

Example 1-10: Emulsion

		%
5	(1) Vitamin A	1.0
	(2) Polyoxyalkylene modified organopolysiloxane ^{*5}	1.0
	(3) Ethanol	2.0
	(4) Glycerol	10.0
10	(5) Sorbitol 70% solu.	3.0
	(6) Propylene glycol	3.0
	(7) Carboxyvinyl polymer	0.3
	(8) KOH	0.1
	(9) Methylparaben	0.1
15	(10) Cetanol	2.5
	(11) Vasoline	2.0
	(12) Squalane	10.0
	(13) Isopropyl myristate	5.0
	(14) Glyceryl monostearate	2.0
20	(15) POE(25 mol) cetyl ether	2.0
	(16) Purified water	Balance

^{*5}: General formula C (Polyoxyethylene group 15 wt%, Polyoxypropylene group 10 wt%, M.W. 5000)

25

(Preparation method)

The present emulsion was obtained in a conventional way. The present emulsion exhibited excellent skin improvement effects.

30

Example 1-11: Emulsion

		%
35	(1) Vitamin A	0.3
	(2) Polyoxyalkylene modified organopolysiloxane ^{*6}	0.2
	(3) Ethanol	5.0
40	(4) Glycerol	5.0
	(5) Sorbitol	2.0
	(6) Propylene glycol	5.0
	(7) Carboxyvinyl polymer	0.2
	(8) KOH	0.06
45	(9) Methyl paraben	0.2
	(10) POE(60 mol) hydrogenated castor oil	1.0
	(11) Squalane	3.0
	(12) Isopropyl myristate	3.0
	(13) Indometacin	0.05
50	(14) Purified water	Balance

^{*6}: General formula D (Polyoxyethylene group 40 wt%, M.W. 7000)

55 (Preparation method)

The present emulsion was obtained in a conventional way. The present emulsion exhibited excellent skin improvement effects.

Example 1-12: Night cream

		%
5	(1) Squalane	20.0
	(2) Liquid paraffin	10.0
	(3) Isopropyl myristate	6.0
	(4) Butyl paraben	0.2
10	(5) Polyoxyalkylene modified organopolysiloxane**7	3.0
	(6) Diglycerol diisostearate	1.0
	(7) Vaseline	4.0
	(8) Solid paraffin	2.0
	(9) Vitamin A	0.3
15	(10) Propylene glycol	4.0
	(11) Glycerol	10.0
	(12) Magnesium sulfate	0.3
	(13) Purified water	Balance

**7: General formula A (Polyoxyethylene group 20 wt%, M.W. 6000)

(Preparation method)

25 The present night cream was obtained in a conventional way. The present night cream exhibited excellent skin improvement effects.

Example 2-1 to 2-5

30 A cream having the following composition was prepared and the skin roughening improvement effects thereof were studied. The vitamin A and the sugars used are shown in Table 2-1 below.

		%
35	(1) Cetostearyl alcohol	3.5
	(2) Squalane	30.0
	(3) Beeswax	3.0
	(4) Reduced lanolin	5.0
	(5) Ethyl paraben	0.3
40	(6) Polyoxyethylene (50 mol) oleyl alcohol ether	2.0
	(7) Stearic monoglyceride	2.0
	(8) Sugar (see Table 2-1)	0.5
	(9) Perfume	0.03
	(10) Vitamin A (see Table 2-1)	0.0001
45	(11) Glycerol	15.0
	(12) Purified water	Balance

(Preparation method)

50 (1), (2), (3), (4), (5), (6), (7), (8), (9) and (10) were heated and dissolved. The resultant solution was maintained at 75 °C, followed by adding thereto, under stirring, (11) and (12) heated to 75 °C. The mixture was stirred and emulsified in a homomixer, followed by cooling to obtain a cream.

Comparative Example 2-1

The same formulation except that the sugar was removed from the formulation of the above Example.

Comparative Example 2-2

The same formulation except that the vitamin A was removed from the formulation of the above Example.

Table 2-1

	Vitamin A	Sugars
Example 2-1	all-trans	Sucrose
Example 2-2	13-cis	Maltitol
Example 2-3	all-trans	D-mannitol
Example 2-4	13-cis	Sorbitol
Example 2-5	all-trans	Fructose
Comp. Ex. 2-1	13-cis	None
Comp. Ex. 2-2	None	Sorbitol

Skin Roughening Improvement Test Method

One hundred subjects having psoriasis-like and skin-roughening-like skin affections were divided into 5 groups as test panels and the creams of Examples 2-1 to 2-5 and Comparative Examples 2-1 to 2-2 were used by one group (20 members) each. That is, Examples 2-1 to 2-5 samples were applied to the left-side faces of the panel twice a day and Comparative Examples 2-1 and 2-2 were applied to the right-side faces of the panel for continuous 3 months. Thereafter, the degree of the overall improvements after the use was visually determined, when compared before the use.

The results are shown in Table 2-2.

Table 2-2: Test Results of Practical Use (Degree of Overall Improvement)

Degree of overall improvement	Example										Comparative Example			
	2-1		2-2		2-3		2-4		2-5		2-1		2-2	
	Psori-asis rough-ening	Skin rough-ening	Psori-asis rough-ening	Skin rough-ening	Psori-asis rough-ening	Skin rough-ening	Psori-asis rough-ening	Skin rough-ening	Psori-asis rough-ening	Skin rough-ening	Psori-asis rough-ening	Skin rough-ening	Psori-asis rough-ening	Skin rough-ening
Remarkable improvement	7	6	6	6	6	6	7	6	6	5	5	0	0	0
Some improvement	2	2	3	2	1	2	2	2	2	3	4	4	5	5
No change	1	2	1	2	3	1	2	2	2	2	19	19	20	20
Change for the worse	0	0	0	0	0	0	0	0	0	0	2	2	0	0
Total (No. of Person)	10	10	10	10	10	10	10	10	10	10	25	25	25	25
Degree of effectiveness %														
Those better than "some improvement"	90	80	90	80	70	90	80	80	80	80	16	16	20	20

As is clear from the results shown in Table 2-2, the products of Examples 2-1 to 2-5 according to the present invention had synergistically excellent skin roughening improvement effects when compared with those of Comparative Examples 2-1 and 2-2.

Example 2-6: Cosmetic lotion

		%
5	(1) Vitamin A	0.1
	(2) D-xylose	0.1
	(3) Glycerol	1.0
	(4) Purified water	Balance
10	(5) Ethanol	7.0
	(6) Polyoxyethylene (20 mol) oleyl alcohol ether	0.5
	(7) Methyl paraben	0.05
	(8) Citric acid	0.01
	(9) Sodium citrate	0.1
15	(10) Camphor	0.01
	(11) Perfume	0.01

(Preparation method)

In the purified water, (2), (3), (8) and (9) were dissolved. Separately, (1), (6), (7) and (11) were dissolved in ethanol and this solution was added to the above purified water to be dissolved, followed by filtration. Thus, the cosmetic lotion was obtained. The cosmetic lotion of the present invention was excellent in the skin roughening improvement effects.

Example 2-7: Pack

		%
30	(1) D-mannose	1.0
	(2) Polyvinyl alcohol	10.0
	(3) Propylene glycol	7.0
	(4) Ethanol	10.0
35	(5) Vitamin A	0.01
	(6) Methyl paraben	0.05
	(7) POE(60 mol) hydrogenated castor oil	0.2
	(8) Perfume	0.05
40	(9) Purified water	Balance

(Preparation method)

In (9), (1), (3) and (6) were added, followed by stirring the mixture to thereby be dissolved. Then, (2) and (7) were added thereto, followed by heating while stirring to dissolve (9) therein. Thereafter, (4) was added, followed by stirring to be dissolved. Thus, the pack was obtained.

The present pack had a synergistically excellent skin roughening improvement effects.

Example 2-8: Compact face powder

		%
5	(1) Vitamin A	0.0005
	(2) Talc	85.4
	(3) Stearic acid	2.5
	(4) Squalane	3.5
10	(5) Sorbitan sesquioleic ester	1.8
	(6) Triethanolamine	1.2
	(7) D-glucosamine	2.5
	(8) Pigment	q.s.
15	(9) Perfume	q.s.

(Preparation method)

20 The talc and the pigment were sufficiently mixed by a kneeder (Powder portion). The triethanolamine was added to 50% corresponding amount of the purified water and the mixture was maintained at 70 °C (Aqueous phase). The components of the present invention other than the perfume were mixed and dissolved under heating at 70 °C (Oil phase). The oil phase was added to the aqueous phase, followed by uniformly emulsified by a homomixer and the resultant emulsified mixture was added to the powder portion, followed by kneeding the same by a kneeder, followed by evaporating the water and by treating the same by a grinder. Furthermore, the perfume was uniformly sprayed and the resultant product was compression molded.

The resultant compact face powder was excellent in the improvement effects to the skin.

Example 2-9: Lipstick

		%
35	(1) Vitamin A	0.00001
	(2) Microcrystalline wax	3.0
	(3) Beeswax	3.0
	(4) Ceresine wax	5.0
	(5) Liquid paraffin	19.0
40	(6) Squalane	20.0
	(7) Carnauba wax	3.0
	(8) Candellira wax	3.0
	(9) D-mannitol	1.0
	(10) Mixed colorant	7.0
45	(11) Dibutyl hydroxytoluene	0.05
	(12) Perfume	q.s.
	(13) Lanolin	Balance

(Preparation method)

The lipstick was obtained in a conventional way. The present lipstick exhibited excellent skin improvement effects.

Example 2-10: Beauty powder

5

		%
(1)	D-mannitol	50.0
(2)	D-sorbitol	45.0
(3)	Vitamin A	0.1
(4)	Talc	4.9

10

(Preparation method)

15

The present beauty powder was obtained in a conventional way. The present beauty powder exhibited excellent skin improvement effects.

Example 2-11: Emulsion

20

		%
(1)	Vitamin A	1.0
(2)	D-erithorlose	2.5
(3)	Ethanol	2.0
(4)	Glycerol	10.0
(5)	Propylene glycol	3.0
(6)	Carboxyvinyl polymer	0.3
(7)	KOH	0.1
(8)	Methyl paraben	0.1
(9)	Cetanol	2.5
(10)	Vaseline	2.0
(11)	Squalane	10.0
(12)	Isopropyl myristate	5.0
(13)	Isopropylene piconol	0.01
(14)	Glyceryl monostearate	2.0
(15)	POE(25 mol) cetyl ether	2.0
(16)	Purified water	Balance

40

(Preparation method)

45

The present emulsion was obtained in a conventional way. The present emulsion exhibited excellent skin improvement effects.

50

55

Example 2-12: Emulsion

5

10

15

20

(Preparation method)

25

Example 2-13: Night cream

30

35

40

45

(Preparation method)

The present night cream was obtained in a conventional way. The present night cream exhibited synergistically excellent skin improvement effects.

50 Examples 3-1 to 3-5

Creams having the following compositions were prepared and the skin roughening improvement effects thereof were studied. The anti-inflammatory agents formulated are listed in Table 3-1.

55

		%
(1)	Vitamin A	0.3
(2)	L-arabinose	2.5
(3)	Ethanol	5.0
(4)	Glycerol	5.0
(5)	Propylene glycol	5.0
(6)	Carboxyvinyl polymer	0.2
(7)	KOH	0.06
(8)	Methyl paraben	0.2
(9)	POE(60 mol) hydrogenated castor oil	1.0
(10)	Squalane	3.0
(11)	Isopropyl myristate	3.0
(12)	Monoammonium glycyrrhizinate	0.05
(13)	Purified water	Balance

		%
(1)	Squalane	18.0
(2)	Liquid paraffin	12.0
(3)	Isopropyl myristate	7.0
(4)	Propyl paraben	0.2
(5)	Diglycerine dioleate	2.0
(6)	POE(5 mol) hydrogenated castor oil	2.0
(7)	Vaseline	6.0
(8)	Vitamin A	0.5
(9)	Erithritol	10.0
(10)	Sucrose	5.0
(11)	Purified water	Balance

		%
(1)	Cetostearyl alcohol	3.5
(2)	Squalane	30.0
(3)	Beeswax	3.0
(4)	Reduced lanolin	5.0
(5)	Ethyl paraben	0.3
(6)	Polyoxyethylene (50 mol) oleyl alcohol ether	2.0
(7)	Stearic monoglyceride	2.0
(8)	Inflammatory agent (see Table 3-1)	0.1
(9)	Perfume	0.03
(10)	Vitamin A	0.0001
(11)	Glycerol	15.0
(12)	Purified water	Balance

(Preparation method)

(1), (2), (3), (4), (5), (6), (7), (8), (9) and (10) were heated and dissolved. The resultant solution was maintained at 75°C, followed by adding thereto, under stirring, (11) and (12) heated to 75°C. The mixture was stirred and emulsified in a homomixer, followed by cooling to obtain a cream.

Comparative Example 3-1

The same formulation except that the anti-inflammatory agents was removed from the formulation of Example 3-1.

Comparative Example 3-2

The same formulation except that the vitamin A was removed from the formulation of Example 3-1.

Table 3-1

	Drugs formulated
Example 3-1	Glycyrrhizic ammonium
Example 3-2	Allantoin
Example 3-3	Glycyrrhetic stearyl
Example 3-4	Hydrocortisone
Example 3-5	Acetaminophen

Skin Roughening Improvement Test Method

One hundred subjects having psoriasis-like and skin-roughening-like skin affections were divided into 5 groups as test panels and the creams of Examples 3-1 to 3-5 and Comparative Examples 3-1 to 3-2 were used by one group (20 members) each. That is, Examples 3-1 to 3-5 samples were applied to the left-side faces of the panel twice a day and Comparative Examples 3-1 and 3-2 were applied to the right-side faces of the panel for continuous 3 months. Thereafter, the degree of the overall improvements after the use was visually determined, when compared before the use.

The results are shown in Table 3-2.

Table 3-2: Test Results of Practical Use (Degree of Overall Improvement)

Degree of overall improvement	Example						Comparative Example					
	3-1	3-2	3-3	3-4	3-5	3-1	3-1	3-1	3-1	3-1	3-2	3-2
	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening	Psori- Skin rough-asis ening
Remarkable improvement	7	6	6	6	6	6	6	6	6	6	6	6
Some improvement	2	2	3	2	2	2	2	2	2	2	2	2
No change	1	2	1	2	2	2	2	2	2	2	2	2
Change for the worse	0	0	0	0	0	0	0	0	0	0	0	0
Total (No. of Person)	10	10	10	10	10	10	10	10	10	10	10	10
Degree of effectiveness %	90	80	90	80	80	80	80	80	80	80	80	80
Those better than "some improvement"	90	80	90	80	80	80	80	80	80	80	80	80

As is clear from the results shown in Table 3-2, the products of the present invention of Examples 3-1 to 3-5 had the synergistically excellent skin roughening improvement effects, when compared with the products of Comparative Examples 3-1 and 3-2.

Example 3-6: Cosmetic lotion

		%
5	(1) Vitamin A	0.00001
	(2) Bethamethasone	0.01
	(3) Glycerol	1.0
	(4) Maltitol	0.3
10	(5) Purified water	Balance
	(6) Ethanol	7.0
	(7) Polyoxyethylene (50 mol) oleyl alcohol ether	1.0
	(8) Methyl paraben	0.05
	(9) Oleyl alcohol	1.0
15	(10) Lactic acid	0.01
	(11) Sodium lactate	0.1
	(12) Perfume	0.01

20 (Preparation method)

In the purified water, (3), (4), (10) and (11) were dissolved. Separately, (1), (2), (7), (8) and (12) were dissolved in ethanol and this solution was added to the above purified water to be dissolved, followed by filtration. Thus, the cosmetic lotion was obtained. The cosmetic lotion of the present invention was excellent in the skin roughening improvement effects.

Example 3-7: Pack

		%
30	(1) Dexamethasone acetate	0.5
	(2) Polyvinyl alcohol	10.0
	(3) Propylene glycol	7.0
35	(4) Ethanol	10.0
	(5) Vitamin A	0.01
	(6) Methyl paraben	0.05
	(7) POE(60 mol) hydrogenated castor oil	0.2
	(8) Perfume	0.05
40	(9) Purified water	Balance

45 (Preparation method)

In (9), (1), (3), (6) and (7) were added, followed by stirring, whereby the mixture was dissolved. Then, (2) was added thereto and the mixture was stirred under heating. Then, (4) containing (9) dissolved therein was added and stirred, whereby the mixture was dissolved to obtain the pack.

The present pack exhibited the excellent skin roughening improvement effects.

50

55

Example 3-8: Compact face powder

		%
5	(1) Vitamin A	0.0005
	(2) Talc	85.4
	(3) Stearic acid	2.5
	(4) Squalane	3.5
10	(5) Sorbitan sesquioleic ester	1.8
	(6) Triethanolamine	1.2
	(7) Diclofenac	0.01
	(8) Pigment	q.s.
15	(9) Perfume	q.s.

(Preparation method)

20 The talc and the pigment were sufficiently mixed by a kneeder (Powder portion). The triethanolamine was added to 50% corresponding amount of the purified water and the mixture was maintained at 70 °C (Aqueous phase). The components of the present invention other than the perfume were mixed and dissolved under heating at 70 °C (Oil phase). The oil phase was added to the aqueous phase, followed by uniformly emulsified by a homomixer and the resultant emulsified mixture was added to the powder portion, followed by kneeding the same by a kneeder, followed by evaporating the water and by treating the same by a grinder. Furthermore, the perfume was uniformly sprayed and the resultant product was compression molded.

The resultant compact face powder was excellent in the improvement effects to the skin.

The present compact face powder exhibited excellent skin improvement effects.

30 Example 3-9: Lipstick

		%
35	(1) Vitamin A	0.00001
	(2) Microcrystalline wax	3.0
	(3) Beeswax	3.0
	(4) Ceresin wax	5.0
40	(5) Liquid paraffin	19.0
	(6) Squalane	20.0
	(7) Carnauba wax	3.0
	(8) Candellira wax	3.0
	(9) Glycyrrhizinic stearyl	5.0
45	(10) Mixed colorant	7.0
	(11) Dibutyl hydroxytoluene	0.05
	(12) Perfume	q.s.
	(13) Lanolin	Balance

50 (Preparation method)

The lipstick was obtained in a conventional way. The present lipstick prevented the formation of roughening on the lip.

Example 3-10: Emulsion

5

10

15

20

		%
(1)	Vitamin A	1.0
(2)	Hydrocortisone acetate	0.05
(3)	Ethanol	2.0
(4)	Glycerol	10.0
(5)	Mannitol	3.0
(6)	Propylene glycol	3.0
(7)	Carboxyvinyl polymer	0.3
(8)	KOH	0.1
(9)	Methyl paraben	0.1
(10)	Cetanol	2.5
(11)	Vaseline	2.0
(12)	Squalane	10.0
(13)	Isopropyl myristate	5.0
(14)	Glyceryl monostearate	2.0
(15)	POE(25 mol) cetyl ether	2.0
(16)	Purified water	Balance

25 (Preparation method)

The present emulsion was obtained in a conventional way. The present emulsion exhibited excellent skin improvement effects.

30 Example 3-11: Emulsion

35

40

45

		%
(1)	Vitamin A	0.3
(2)	Indometacin	0.3
(3)	Ethanol	5.0
(4)	Glycerol	5.0
(5)	Propylene glycol	5.0
(6)	Carboxyvinyl polymer	0.2
(7)	KOH	0.06
(8)	Methyl paraben	0.2
(9)	POE(60 mol) hydrogenated castor oil	1.0
(10)	Squalane	3.0
(11)	Isopropyl myristate	3.0
(12)	Purified water	Balance

50 (Preparation method)

The present emulsion was obtained in a conventional way. The present emulsion exhibited excellent skin improvement effects.

55

Example 3-12: Night cream

		%
5	(1) Squalane	10.0
	(2) Liquid paraffin	10.0
	(3) Vaseline	3.0
	(4) Cetyl octanoate	10.0
10	(5) Dibutyl phthalate	5.0
	(6) Glycyrrhizinic stearyl	0.1
	(7) Indometacin	0.2
	(8) Butyl paraben	0.2
	(9) Diglycerine triisostearate	2.0
15	(10) Diglycerine monoisostearate	1.5
	(11) Vitamin A	0.1
	(12) Glycerol	10.0
	(13) Propylene glycol	6.0
20	(14) Purified water	Balance

(Preparation method)


25 The present night cream was obtained in a conventional way. The present night cream exhibited excellent skin improvement effects.

[Industrial Applicability]

30 The external skin treatment composition according to the present invention are useful as an external skin treatment composition capable of preventing the epidermal disabilities and synergistically of improving the changes and disabilities due to aged skins or sunlight exposure, with taking safety into consideration.

Claims

- 35 1. An external skin treatment composition comprising (i) vitamin A and (ii) at least one skin roughening improvement aid selected from the group consisting of (a) polyoxyalkylene modified organopolysiloxanes, (b) sugars, and (c) anti-inflammatory drugs.
- 40 2. A composition according to claim 1, wherein the skin roughening improvement aid is at least one polyoxyalkylene modified organopolysiloxane selected from the group consisting of the compounds represented by the formulae (A), (B), (C) and (D), wherein R represents an alkyl group having 1 to 3 carbon atoms or a phenyl group, R' is hydrogen or an alkyl group having 1 to 12 carbon atoms, p is an integer of 1 to 5, m is an integer of 5 to 100, n and x are an integer of 1 to 50, and t and y are an integer of 0 to 50.
- 45
- 50
- 55



3. A composition according to claim 2, wherein the amount of the vitamin A formulated in the composition is 0.00001 to 5.0% by weight and the amount of the polyoxyalkylene modified organopolysiloxane formulated in the composition is 0.1 to 20% by weight.

4. A composition according to claim 1, wherein the skin roughening improvement aid is at least one sugar selected from the group consisting of monosaccharides, oligosaccharides and sugar alcohols.
5. A composition according to claim 4, wherein the amount of vitamin A formulated in the composition is 0.00001 to 5.0% by weight and the amount of the sugar formulated in the composition is 0.1% by weight or more.
6. A composition according to claim 1, wherein the skin roughening improvement aid is at least one anti-inflammatory agent selected from the group consisting of hydrocortisone, hydrocortisone acetate, prednisolone, methyl prednisolone, prednisolone acetate, prednisolone acetate propionate, dexamethasone, betamethasone, triamcinolone, dexamethasone acetate, betamethasone valerate, triamcinolone acetate, aspirin salicylic acid, acetaminophen, methyl salicylate, glycol salicylate, mefenamic acid, flufenamic acid, indometacin, diclofenac, ketoprofen, ibuprofen, flurbiprofen, fenbufen, lufexamac, piroxicam, oxyphenbutazone, mepirizole, ibuprofen piconol, clidanac, phenylbutazone, naproxen, glycyrrhetin, glycyrrhizin, glycyrrhetic acid and the salts and esters thereof, glycyrrhizic acid and the salts and esters thereof, azulene, camphor, thymol, allantoin, etc.
7. A composition according to claim 6, wherein the amount of vitamin A formulated in the composition is 0.00001 to 5.0% by weight and the amount of the anti-inflammatory agent formulated in the composition is 0.0001 to 5.0% by weight.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP93/00968

A. CLASSIFICATION OF SUBJECT MATTER

Int. Cl⁵ A61K7/48, A61K7/00, A61K31/57, A61K31/19

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int. Cl⁵ A61K7/48, A61K7/00, A61K31/57, A61K31/19

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP, A, 273202 (VANSCOTT E. J.), July 6, 1988 (06. 07. 88), & JP, A, 63-166837	1, 6, 7
X	JP, A, 1-186811 (Sunstar K.K.), July 26, 1989 (26. 07. 89), (Family: none)	1, 6, 7
X	DE, A, 3327840 (BLENDAX WERKE SCHNEIDER), September 20, 1984 (20. 09. 84), & JP, A, 60-69012 & EP, A, 155344 & US, A, 4743442	1, 4, 5
X	JP, A, 59-95210 (SEWA K.K.), June 1, 1984 (01. 06. 84), (Family: none)	1, 4, 5
X	JP, A, 57-131716 (SEWA K.K.), August 14, 1982 (14. 08. 82), (Family: none)	1, 4, 5

☒ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

September 27, 1993 (27. 09. 93)

Date of mailing of the international search report

October 12, 1993 (12. 10. 93)

Name and mailing address of the ISA/

Japanese Patent Office

Facsimile No.

Authorized officer

Telephone No.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP93/00968

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP, A, 58-140008 (SEWA K.K.), August 19, 1983 (19. 08. 83), (Family: none)	1, 4, 5
X	JP, A, 1-186809 (SUNSTAR K.K.), July 26, 1989 (26. 07. 89), (Family: none)	1, 6, 7
X	EP, A, 158090 (ISMAIL R), October 16, 1985 (16. 10. 85), & DE, A, 3410641 & JP, A, 61-40210 & US, A, 4938960	1, 6, 7
Y	JP, A, 63-10710 (KAO CORP.), January 18, 1988 (18. 01. 88), (Family: none)	1-3
Y	EP, A, 255364 (AVON PRODUCTS INC.), February 3, 1988 (03. 02. 88), & JP, A, 63-35510 & US, A, 4888363	1-3
Y	WO, A, 8606275 (AVON PRODUCTS INC.), November 6, 1986 (06. 11. 86), & DE, A, 3684417 & EP, A, 224504 & JP, A, 62-502546	1-3
Y	DE, A, 3431755 (L'OREAL SA), March 14, 1985 (14. 03. 85), & FR, A, 2550940 & JP, A, 60-78914 & US, A, 4608392	1, 6, 7
Y	WO, A, 8503434 (NEUTROGENA CORP.), August 15, 1985 (15. 08. 85), & EP, A, 172228 & JP, A, 61-501091 & US, A, 4678663	1, 6, 7
A	JP, A, 1-63031 (SHISEIDO K.K.), March 9, 1989 (09. 03. 89), (Family: none)	1-3